

Body Mass Index Profiles of Police Cadets Belonging to Three Different Academy Classes

Connelly, Greta; Johnson, Quincy; Diehl, Cody; Kornhauser, Charlie; Lockie, Robert G.; Orr, Rob Marc; Dawes, Jay

Licence:
CC BY-NC-ND

[Link to output in Bond University research repository.](#)

Recommended citation(APA):

Connelly, G., Johnson, Q., Diehl, C., Kornhauser, C., Lockie, R. G., Orr, R. M., & Dawes, J. (2019). *Body Mass Index Profiles of Police Cadets Belonging to Three Different Academy Classes*. Poster session presented at American College of Sports Medicine Central States Annual Meeting, Broken Arrow, Oklahoma, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.



BODY MASS INDEX PROFILES OF POLICE CADETS BELONGING TO THREE DIFFERENT ACADEMY CLASSES

¹Greta Connelly, ¹Quincy R. Johnson, ¹Cody L. Diehl, ²Charles L. Kornhauser, ³Robert G. Lockie, ^{4,5}Robin M. Orr, J. ¹Jay Dawes

¹ Department of Health and Human Performance, Oklahoma State University, Stillwater, OK, USA; ² Colorado State Patrol Training Academy, Golden, CO, USA, QLD, AUS; ³ California State University Fullerton, Fullerton, CA, USA; ⁴ Tactical Research Unit, Robina, QLD, ASU



ABSTRACT

Body mass index (BMI) is a method frequently utilized for assessing an individual's health status, and for predicting morbidity and mortality risk. Although BMI has been criticized as being inaccurate for individuals with larger body frames and greater muscle mass, it does provide an easy and cost-effective health risk assessment tool that can be used to broadly determine if more invasive measures of body composition are warranted.

Purpose: To assess the general health status of police cadets attending three separate training academies from one US based law enforcement agency.

Methods: Archived data for police cadets (n = 127) from one US law enforcement agency was provided to the primary investigator for analysis. BMI was then calculated using the Quetelet Index Formula ($BMI = \frac{kg}{m^2}$).

Results: Surprisingly, 2% of all cadets were classified as underweight, 33% as normal weight, 44% as overweight, and 22% as obese.

Conclusions: Based on these findings, it appears that the majority of police cadets attending training academy would be classified as overweight or obese. These results can be used to inform cadets about their general health status and guide training programs to reduce the risk of premature morbidity and mortality in this population.

INTRODUCTION

Body mass index (BMI) can be used to classify individuals into different categories of health. Based on their BMI score an individual may be classified into one of six major categories:

- Underweight ($BMI < 18.5 \text{ kg/m}^2$)
- Normal weight ($18.5 < 25 \text{ kg/m}^2$)
- Overweight ($25 < 30 \text{ kg/m}^2$)
- Low obesity ($30 < 35 \text{ kg/m}^2$)
- Medium obesity ($35 < 40 \text{ kg/m}^2$)
- Extremely obese ($> 40 \text{ kg/m}^2$)

Individuals that have a BMI of 25 or higher, are shown to be at higher risk for certain medical complications, such as heart disease, hypertension, type 2 diabetes, gallstones, and some types of cancers (1,2,3).

Even though BMI has been critiqued for being inaccurate for some individuals with larger body frames, and greater muscle mass, it does provide an easy and cost effective health risk assessment.




METHODS

Archived data for 127 (M=111, F=16 age = 31.3 ± 7.3 yrs ; Ht= 178.4 ± 8.9 cm; Wt = 87.5 ± 16.7 kg; BMI = $27.4 \pm 4.6 \text{ kg/m}^2$) police cadets was utilized for this analysis. Prior to academy training, cadets had their height and weight measured via a doctor's beam scale in order to track changes in body mass during academy training and to assess general health status. BMI was calculated by dividing body mass in kilograms (kg) by height in meters squared (m^2). Based on their BMI score, individuals were classified into one of four major categories: underweight ($BMI < 18.5 \text{ kg/m}^2$), normal weight ($BMI = 18.5 - 24.9 \text{ kg/m}^2$), overweight ($BMI = 25 - 29.9 \text{ kg/m}^2$) obesity ($BMI \geq 30 \text{ kg/m}^2$).


RESULTS

Figure 1. BMI classifications



RESULTS (CONT.)

Figure 2. Average BMI classifications for cadets by year



CONCLUSIONS

Based on these findings, it appears that the majority of police cadets attending training academy would be classified as overweight or obese. These findings are concerning based on the known health risk associated with failure to attain and maintain an ideal body mass level. BMI may provide a simple measurement of preliminary health status for cadets participating in academy training that can later be used for providing physical activity and nutritional recommendations.

REFERENCES

1. Charles, L. E., Burchfiel, C. M., Violanti, J. M., Fekedulegn, D., Slaven, J. E., Browne, R. W., ... Andrew, M. E. (2008). Adiposity Measures and Oxidative Stress Among Police Officers. *Obesity*, 16(11), 2489–2497. doi: 10.1038/oby.2008.395
2. Pasco, J. A., Holloway, K. L., Dobbins, A. G., Kotowicz, M. A., Williams, L. J., & Brennan, S. L. (2014). Body mass index and measures of body fat for defining obesity and underweight: a cross-sectional, population-based study. *BMC Obesity*, 1(1). doi: 10.1186/2052-9538-1-9
3. Stommel, M., & Schoenborn, C. A. (2009). Accuracy and usefulness of BMI measures based on self-reported weight and height: findings from the NHANES & NHIS 2001-2006. *BMC Public Health*, 9(1). doi: 10.1186/1471-2458-9-421